# BRIGHT FUTURES HANDOUT ▶ PARENT

# 2 YEAR VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.





# **HOW YOUR FAMILY IS DOING**

- Take time for yourself and your partner.
- Stay in touch with friends.
- Make time for family activities. Spend time with each child.
- Teach your child not to hit, bite, or hurt other people. Be a role model.
- If you feel unsafe in your home or have been hurt by someone, let us know.
  Hotlines and community resources can also provide confidential help.
- Don't smoke or use e-cigarettes. Keep your home and car smoke-free.
  Tobacco-free spaces keep children healthy.
- Don't use alcohol or drugs.
- Accept help from family and friends.
- If you are worried about your living or food situation, reach out for help.
  Community agencies and programs such as WIC and SNAP can provide information and assistance.



# **TALKING AND YOUR CHILD**

- Use clear, simple language with your child. Don't use baby talk.
- Talk slowly and remember that it may take a while for your child to respond. Your child should be able to follow simple instructions.
- Read to your child every day. Your child may love hearing the same story over and over.
- Talk about and describe pictures in books.
- Talk about the things you see and hear when you are together.
- Ask your child to point to things as you read.
- Stop a story to let your child make an animal sound or finish a part of the story.



### YOUR CHILD'S BEHAVIOR

- Praise your child when he does what you ask him to do.
- Listen to and respect your child. Expect others to as well.
- Help your child talk about his feelings.
- Watch how he responds to new people or situations.
- Read, talk, sing, and explore together. These activities are the best ways to help toddlers learn.
- Limit TV, tablet, or smartphone use to no more than 1 hour of high-quality programs each day.
  - It is better for toddlers to play than to watch TV.
  - Encourage your child to play for up to 60 minutes a day.
- Avoid TV during meals. Talk together instead.



# **TOILET TRAINING**

- Begin toilet training when your child is ready.
  Signs of being ready for toilet training include
  - Staying dry for 2 hours
  - Knowing if she is wet or dry
  - Can pull pants down and up
  - Wanting to learn
  - Can tell you if she is going to have a bowel movement
- Plan for toilet breaks often. Children use the toilet as many as 10 times each day.
- Teach your child to wash her hands after using the toilet.
- Clean potty-chairs after every use.
- Take the child to choose underwear when she feels ready to do so.

Helpful Resources: National Domestic Violence Hotline: 800-799-7233 | Smoking Quit Line: 800-784-8669 Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

### **SAFETY**

- Make sure your child's car safety seat is rear facing until he reaches the highest weight or height allowed by the car safety seat's manufacturer. Once your child reaches these limits, it is time to switch the seat to the forwardfacing position.
- Make sure the car safety seat is installed correctly in the back seat. The harness straps should be snug against your child's chest.
- Children watch what you do. Everyone should wear a lap and shoulder seat belt in the car.
- Never leave your child alone in your home or yard, especially near cars or machinery, without a responsible adult in charge.
- When backing out of the garage or driving in the driveway, have another adult hold your child a safe distance away so he is not in the path of
- Have your child wear a helmet that fits properly when riding bikes and trikes.
- If it is necessary to keep a gun in your home, store it unloaded and locked with the ammunition locked separately.

# WHAT TO EXPECT AT YOUR CHILD'S 2½ YEAR VISIT


### We will talk about

- Creating family routines
- Supporting your talking child
- Getting along with other children
- Getting ready for preschool
- Keeping your child safe at home, outside, and in the car

Consistent with Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th Edition

For more information, go to https://brightfutures.aap.org.

American Academy of Pediatrics DEDICATED TO THE HEALTH OF ALL CHILDREN®



The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

# ownloaded from http://publications.aap.org/pati<del>bytEtlysticiter.pdf2601/f18df36201/ppanydosum</del>ent194/1200165/peo\_document194\_en.pd

# **Hepatitis A Vaccine:**

# What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

# 1. Why get vaccinated?

Hepatitis A vaccine can prevent hepatitis A.

**Hepatitis A** is a serious liver disease. It is usually spread through close, personal contact with an infected person or when a person unknowingly ingests the virus from objects, food, or drinks that are contaminated by small amounts of stool (poop) from an infected person.

Most adults with hepatitis A have symptoms, including fatigue, low appetite, stomach pain, nausea, and jaundice (yellow skin or eyes, dark urine, light-colored bowel movements). Most children less than 6 years of age do not have symptoms.

A person infected with hepatitis A can transmit the disease to other people even if he or she does not have any symptoms of the disease.

Most people who get hepatitis A feel sick for several weeks, but they usually recover completely and do not have lasting liver damage. In rare cases, hepatitis A can cause liver failure and death; this is more common in people older than 50 years and in people with other liver diseases.

Hepatitis A vaccine has made this disease much less common in the United States. However, outbreaks of hepatitis A among unvaccinated people still happen.

# 2. Hepatitis A vaccine

Children need 2 doses of hepatitis A vaccine:

- First dose: 12 through 23 months of age
- Second dose: at least 6 months after the first dose

Infants 6 through 11 months old traveling outside the United States when protection against hepatitis A is recommended should receive 1 dose of hepatitis A vaccine. These children should still get 2 additional doses at the recommended ages for long-lasting protection.

**Older children and adolescents** 2 through 18 years of age who were not vaccinated previously should be vaccinated.

**Adults** who were not vaccinated previously and want to be protected against hepatitis A can also get the vaccine.

Hepatitis A vaccine is also recommended for the following people:

- International travelers
- Men who have sexual contact with other men
- People who use injection or non-injection drugs
- People who have occupational risk for infection
- People who anticipate close contact with an international adoptee
- People experiencing homelessness
- People with HIV
- People with chronic liver disease

In addition, a person who has not previously received hepatitis A vaccine and who has direct contact with someone with hepatitis A should get hepatitis A vaccine as soon as possible and within 2 weeks after exposure.

Hepatitis A vaccine may be given at the same time as other vaccines.



# 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

 Has had an allergic reaction after a previous dose of hepatitis A vaccine, or has any severe, lifethreatening allergies

In some cases, your health care provider may decide to postpone hepatitis A vaccination until a future visit.

Pregnant or breastfeeding people should be vaccinated if they are at risk for getting hepatitis A. Pregnancy or breastfeeding are not reasons to avoid hepatitis A vaccination.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting hepatitis A vaccine.

Your health care provider can give you more information.

# 4. Risks of a vaccine reaction

 Soreness or redness where the shot is given, fever, headache, tiredness, or loss of appetite can happen after hepatitis A vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

# 5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at <a href="https://www.vaers.hhs.gov">www.vaers.hhs.gov</a> or call 1-800-822-7967. VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.

# 6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at <a href="https://www.hrsa.gov/vaccinecompensation">www.hrsa.gov/vaccinecompensation</a> or call 1-800-338-2382 to learn about the program and about filing a claim.

# 7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at <a href="www.fda.gov/">www.fda.gov/</a> <a href="www.fda.gov/">waccines-blood-biologics/vaccines</a>.
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call 1-800-232-4636 (1-800-CDC-INFO) or
- Visit CDC's website at www.cdc.gov/vaccines.



