

Acetaminophen (Tylenol®) Dosage Chart for Infants and Children

Dosing Instructions: Give every 4–6 hours as needed for fever or pain. Do NOT give more than 5 doses in 24 hours. Do NOT use with any other medicine containing acetaminophen. If possible, use weight to dose; otherwise, use age. *Talk with your child's doctor if you have a dosing question or your child does not fit into the weight/age range listed.*

Age	0–3 months	4–11 months	12–23 months	2–3 years	4–5 years	6–8 years	9–10 years	11 years	12+ years
Weight (pounds)	6.6–11 lbs	11+–17 lbs	17+–23 lbs	23+–35 lbs	35+–46 lbs	46+–60 lbs	60+–70 lbs	70+–95 lbs	95+ lbs
Infants Acetaminophen Liquid (160 mg / 5 mL)	1.25 mL (40 mg)	2.5 mL (80 mg)	3.75 mL (120 mg)	—	—	—	—	—	—
Acetaminophen Liquid (160 mg / 5 mL)	—	—	—	5 mL (160 mg)	7.5 mL (240 mg)	10 mL (320 mg)	12.5 mL (400 mg)	15 mL (480 mg)	20 mL (640 mg)
Acetaminophen Chewables (160 mg)	—	—	—	1 tablet	1½ tablets	2 tablets	2½ tablets	3 tablets	4 tablets
Acetaminophen Tablets (325 mg)	—	—	—	—	—	1 tablet	1 tablet	1 tablets	2 tablets

Liquid Medicines: Use the syringe or dosing cup that comes with the medication to give an accurate dose. Kitchen teaspoons or tablespoons should not be used.

See reverse side for ibuprofen dosing table. Updated October 2024.

BRV1289328 KN2028



Ibuprofen (Motrin®, Advil®) Dosage Chart for Infants and Children

Dosing Instructions: Give every 6–8 hours as needed for fever or pain. Do NOT give more than 4 doses in 24 hours. Do NOT use with any other medicine containing ibuprofen. If possible, use weight to dose; otherwise, use age.

Talk with your child's doctor if you have a dosing question or your child does not fit into the weight/age range listed.

Age	6–11 months	12–23 months	2–3 years	4–5 years	6–8 years	9–10 years	11 years	12+ years
Weight (pounds)	11*–17 lbs	17*–23 lbs	23*–35 lbs	35*–46 lbs	46*–60 lbs	60*–70 lbs	70*–95 lbs	95+ lbs
Infant's Ibuprofen Drops (50 mg / 1.25 mL)	1.25 mL (50 mg)	2 mL (80 mg)	2.5 mL (100 mg)	—	—	—	—	—
Liquid Ibuprofen (100 mg / 5 mL)	2.5 mL (50 mg)	4 mL (80 mg)	5 mL (100 mg)	7.5 mL (150 mg)	10 mL (200 mg)	12.5 mL (250 mg)	15 mL (300 mg)	20 mL (400 mg)
Ibuprofen Chewable Tablets (100 mg)	—	—	1 tablet	1½ tablets	2 tablets	2½ tablets	3 tablets	4 tablets
Ibuprofen Tablets (200 mg)	—	—	—	—	1 tablet	1 tablet	1½ tablets	2 tablets

Age Limits: Do NOT use ibuprofen for infants less than 6 months old unless your child's doctor tells you.

Liquid Medicines: Use the syringe or dosing cup that comes with the medication to give an accurate dose. Kitchen teaspoons or tablespoons should not be used.

See reverse side for acetaminophen dosing table. Updated October 2024.
BRV1289328 KN2028

